

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representation of
The original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

THIS PAGE BLANK (USPTO)



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification : C12N 15/82, 15/52, A01H 5/00	A1	(11) International Publication Number: WO 92/19747 (43) International Publication Date: 12 November 1992 (12.11.92)
<p>(21) International Application Number: PCT/GB92/00742</p> <p>(22) International Filing Date: 23 April 1992 (23.04.92)</p> <p>(30) Priority data: 9108756.9 24 April 1991 (24.04.91) GB</p> <p>(71) Applicant (for all designated States except US): IMPERIAL CHEMICAL INDUSTRIES PLC [GB/GB]; Imperial Chemical House, Millbank, London SW1P 3JF (GB).</p> <p>(72) Inventors; and (75) Inventors/Applicants (for US only) : BRIGHT, Simon, William, Jonathan [GB/GB]; 24 Pound Lane, Marlow, Bucks SR7 2AY (GB). BYROM, David [GB/GB]; 27 Moor Park, Nunthorpe, Middlesbrough, Cleveland TS7 0JJ (GB). FENTEM, Philip, Anthony [GB/GB]; 9 Aggisters Lane, Wokingham, Berkshire RG11 4DW (GB).</p> <p>(74) Agent: HUSKISSON, Frank, Mackie; Imperial Chemical Industries plc, ICI Group Legal Department, Patents, PO Box No 6, Bessemer Road, Welwyn Garden City, Herts AL7 1HD (GB).</p>		<p>(81) Designated States: AT (European patent), AU, BB, BE (European patent), BF (OAPI patent), BG, BJ (OAPI patent), BR, CA, CF (OAPI patent), CG (OAPI patent), CH, CH (European patent), CI (OAPI patent), CM (OAPI patent), CS, DE (European patent), DK (European patent), ES (European patent), FI, FR (European patent), GA (OAPI patent), GB (European patent), GN (OAPI patent), GR (European patent), HU, IT (European patent), JP, KP, KR, LK, LU (European patent), MC (European patent), MG, ML (OAPI patent), MN, MR (OAPI patent), MW, NL (European patent), NO, PL, RO, RU, SE, SE (European patent), SN (OAPI patent), TD (OAPI patent), TG (OAPI patent), US.</p> <p>Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i></p>
(54) Title: PRODUCTION OF POLYHYDROXYALKANOATE IN PLANTS		
<p>(57) Abstract</p> <p>A plant which produces polyhydroxyalkanoate polymer has a recombinant genome which contains one or more than one of the genes specifying enzymes critical to the polyhydroxyalkanoate biosynthetic pathway which occurs in certain micro-organisms such as <i>Alcaligenes eutrophus</i> which naturally produce same. The plant species is preferably an oil-producing plant.</p> <div data-bbox="860 1134 1445 1806" data-label="Chemical-Block"> </div>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	FI	Finland	MI	Mali
AU	Australia	FR	France	MN	Mongolia
BB	Barbados	GA	Gabon	MR	Mauritania
BE	Belgium	GB	United Kingdom	MW	Malawi
BF	Burkina Faso	GN	Guinea	NL	Netherlands
BG	Bulgaria	GR	Greece	NO	Norway
BJ	Benin	HU	Hungary	PL	Poland
BR	Brazil	IE	Ireland	RO	Romania
CA	Canada	IT	Italy	RU	Russian Federation
CF	Central African Republic	JP	Japan	SD	Sudan
CG	Congo	KP	Democratic People's Republic of Korea	SE	Sweden
CH	Switzerland	KR	Republic of Korea	SN	Senegal
CI	Côte d'Ivoire	LI	Liechtenstein	SU	Soviet Union
CM	Cameroon	LK	Sri Lanka	TD	Chad
CS	Czechoslovakia	LU	Luxembourg	TC	Togo
DE	Germany	MC	Monaco	US	United States of America
DK	Denmark	MG	Madagascar		
ES	Spain				

PRODUCTION OF POLYHYDROXYALKANOATE IN PLANTS

This invention relates to the production of polyhydroxyalkanoate in plants.

Poly-3-hydroxybutyrate (PHB) is a linear polyester of D(-)-3-hydroxybutyrate. It was first
5 discovered in Bacillus megaterium in 1925. Polyhydroxybutyrate accumulates in intracellular granules of a wide variety of bacteria. The granules appear to be membrane bound and can be stained with Sudan Black dye. The polymer is
10 produced under conditions of nutrient limitation and acts as a reserve of carbon and energy. The molecular weight of the polyhydroxybutyrate varies from around 50,000 to greater than 1,000,000, depending on the micro-organisms involved, the
15 conditions of growth, and the method employed for extraction of the polyhydroxybutyrate. Polyhydroxybutyrate is an ideal carbon reserve as it exists in the cell in a highly reduced state, (it is virtually insoluble), and exerts negligible
20 osmotic pressure.

Polyhydroxybutyrate and related polyhydroxyalkanoates, such as poly-3-hydroxyvalerate and poly-3-hydroxyoctanoate, are biodegradable thermoplastics of considerable commercial importance.

25 The terms "polyhydroxyalkanoate" and "PHA" as used hereinafter include polymers of 3-hydroxybutyrate, polymers of related hydroxyalkanoates such as 3-hydroxyvalerate, 3-hydroxyhexanoate, 3-hydroxyoctanoate,
30 3-hydroxydecanoate, and also copolymers and

- 2 -

mixtures of more than one of these hydroxy-alkanoates.

5 Polyhydroxyalkanoate is biodegradable and is broken down rapidly by soil micro-organisms. It is thermoplastic (it melts at 180°C) and can readily be moulded into diverse forms using technology well-established for the other thermoplastics materials such as high-density polyethylene which melts at around the same temperature (190°C). The material is ideal for the production of biodegradable packaging which will degrade in landfill sites and sewage farms. The polymer is biocompatible, as well as biodegradable, and is well tolerated by the mammalian, including human, body; its degradation product, 3-hydroxybutyrate, is a normal mammalian metabolite. Polyhydroxybutyrate degrades only slowly in the body making it suitable for medical applications where long term degradation is required.

20 Polyhydroxyalkanoate, produced by the micro-organism Alcaligenes eutrophus, is manufactured, as a copolymer of polyhydroxybutyrate and polyhydroxyvalerate, by Imperial Chemical Industries PLC and sold under the Trade Mark BIOPOL. The nature of the polymer, for example the proportions of PHB and PHV is determined by the substrate supplied in the fermentation. It is normally supplied in the form of pellets for thermoprocessing. However, polyhydroxyalkanoate is more expensive to manufacture by existing methods than, say, polyethylene. It is, therefore, desirable that new, more economic production of polyhydroxyalkanoate be provided.

An object of the present invention is to provide materials and a method for the efficient production of polyhydroxyalkanoate.

5 According to the present invention there is provided a plant adapted for the production of polyhydroxyalkanoate comprising a recombinant genome of an oil-producing plant, which genome contains genes encoding enzymes necessary for catalysing the production of polyhydroxy-alkanoate
10 together with gene regulatory sequences directing expression of the said genes to target plant cell components.

These regulatory sequences include promoter sequences directing expression of the biosynthetic
15 pathway specifically to the developing seed, and transit peptide sequences targeting the enzymes to appropriate subcellular compartments.

The genes encoding the enzyme or enzymes necessary for the catalysis of polyhydroxyalkanoate
20 production may be isolated from a micro-organism, such as Alcaligenes eutrophus, which is known to produce polyhydroxybutyrate and other polyhydroxy-alkanoates.

It is preferable, for reasons which will later
25 be explained, that the plant be of a species which produces substantial quantities of oil, rather than starch. Such plant species are well known and are simply referred to as "oil-seed" crops and include, oilseed rape, canola, soya and sunflower. Methods
30 for the genetic transformation of many oil crops are known; for example, transformation by Agrobacterium tumefaciens methods are suitable for most. Such methods are well-described in the literature and well-known and extensively practised

in the art.

The biosynthesis of polyhydroxybutyrate from the substrate, acetyl-CoA involves three enzyme-catalysed steps, illustrated in Figure 1 herewith.

5 The three enzymes involved are β -ketothiolase, NADP linked acetoacetyl-CoA reductase, and polyhydroxybutyrate synthase, the genes for which have been cloned from Alcaligenes eutrophus (Schubert et al, 1988, J Bacteriol, 170).
10 When cloned into Escherichia coli the three genes are known to facilitate production of polyhydroxyalkanoate up to 30% of the cell weight.

Genes specifying the production of alkanoates higher than the butyrate are known to exist in
15 bacteria. Isolation of the appropriate genes allows expression of these higher polyhydroxy-alkanoates. For example, genes specifying production of the polyhydroxy-octanoate and the -
20 Pseudomonas oleovorans and Pseudomonas eruginosa. However genes for analogous polymers are widespread in bacterial species.

All the microorganisms required for performance of this invention are publicly
25 available from public culture collections.

An important preferred feature of this invention is the use of an oilseed plant for expression of the polyhydroxyalkanoate. The reason behind our selection of oil-producing crops is that
30 such plants naturally produce large amounts of acetyl-CoA substrate (under aerobic conditions) in the developing seed, which is normally used in fatty acid synthesis. Diversion of this substrate into polyhydroxyalkanoate production will reduce

the amount of oil stored by the seed but will have minimal influence on other aspects of the cell's metabolism. It is therefore possible to produce commercially viable quantities of polyhydroxy-
5 alkanoate such as polyhydroxybutyrate in an oilseed.

It has been previously suggested that Alcaligenes eutrophus genes could be expressed in a starch crop but this has certain problems. In
10 order to optimise polyhydroxyalkanoate production in such a crop, it would probably be necessary to down-regulate starch synthesis. However, even if this down-regulation were to be effected it would not guarantee an increased rate of acetyl-CoA
15 production. Moreover, even if this increased production were actually achieved, it is possible that the acetyl-CoA would be rapidly utilised by respiration in the starch crop.

For expression in higher plants the bacterial
20 (for example Alcaligenes eutrophus) genes require suitable promoter and terminator sequences. Various promoters/terminators are available for use. For constitutive expression the cauliflower mosaic virus CaMV35S promoter and nos terminator
25 may be used. It is however preferred to target synthesis of polyhydroxyalkanoate only to the developing oil storage organ of the oilseed such as the embryo of oilseed rape. The promoter of the rape seed storage protein, napin, could be used to
30 obtain embryo specific expression of polyhydroxyalkanoate genes. Expression of the polyhydroxyalkanoate genes during the precise period when lipid is being made will ensure effective competition by the polyhydroxyalkanoate

enzymes for available acetyl-CoA. The promoters of fatty acid synthesis genes whose expressions are switched on at this time are thus most appropriate candidates to be used as polyhydroxyalkanoate gene promoters. Examples of such promoters are those of seed specific isoforms of rapeseed acyl carrier protein (ACP) or β -ketoacyl ACP reductase.

In inserting the polyhydroxyalkanoate genes into eukaryotic cells, consideration has to be given to the most appropriate subcellular compartment in which to locate the enzymes. Two factors are important: the site of production of the acetyl-CoA substrate, and the available space for storage of the polyhydroxyalkanoate polymer.

The acetyl-CoA required for fatty acid synthesis in, for example, developing rapeseed embryo is produced by two routes. The first, direct, route involves the activity of a pyruvate dehydrogenase enzyme located in the plastid. The second route involves the initial production of acetyl-CoA by mitochondrial pyruvate dehydrogenase, lysis to free acetate, and diffusion of the acetate into the plastid where it is re-esterified to CoA by acetyl-CoA synthase. Rapeseed also produces acetyl-CoA in the cytosol, though at a lower rate than in the plastid, via the activity of a cytosolic citrate lyase enzyme.

Considering substrate supply, the bacterial (for example, Alcaligenes) β -ketothiolase enzyme may function in the mitochondrion, using acetyl-CoA produced in excess of the requirements of respiration, or in the cytosol. The regulatory sequences of the invention may thus direct expression of the β -ketothiolase gene to the

- 7 -

mitochondrion or to the cytosol. It is however preferred to target this enzyme to the plastids, where highest rates of acetyl-CoA generation occur.

The mitochondrion lacks sufficient space for storage of the polyhydroxyalkanoate polymer. Significant storage space exists in the plastids, at least in rape embryo. Highest storage space exists in the cytosol, the compartment normally occupied by the oil bodies.

It is not known whether the acetoacetyl-CoA or hydroxybutyryl-CoA pathway intermediates can be transported from plastid to cytosol. Certainly they would not be able to traverse the plastid envelope membrane as CoA esters. Export would require that the acetoacetate or hydroxybutyrate groups are recognised by the transport systems involved in export of fatty acids from plastids. These have been suggested to involve: lysis of the CoA ester, export of the free acid, and resynthesis of the CoA ester in the cytosol; or transfer of the acyl groups to carnitine, and export of acyl carnitine. If acetoacetyl groups may be exported from the plastid by one of these mechanisms then it would be possible to target β -ketothiolase to the plastid, to utilise acetyl-CoA destined for lipid synthesis, and target acetoacetyl-CoA reductase and polyhydroxybutyrate synthase to the cytosol to achieve polymer synthesis in this more spacious compartment. If neither acetoacetate nor hydroxybutyrate groups may be exported from the plastid, polyhydroxyalkanoate synthesis will require that all three pathway enzymes are targeted to this organelle so that they are expressed in the same cell compartment.

To target the three bacterial (such as Alcaligenes eutrophus) enzymes for polyhydroxyalkanoate synthesis to the plant plastid requires the use of specific targeting regulatory elements called transit peptides.

5 Possible sources of plastid stroma targeting sequences are the genes for:

- (a) ribulose biphosphate carboxylase/oxygenase small subunit (RUBISCO ssu);
- 10 (b) acyl carrier protein (ACP);
- (c) β -ketoacyl ACP reductase;
- (d) enolpyruvylshikimate-3-phosphate synthase (EPSPS);
- (e) fructose 1,6-bisphosphatase.

15 Of these the RUBISCO small subunit transit peptide has been shown to direct polypeptides to plastids in both photosynthetic and non-photosynthetic tissues. ACP and β -ketoacyl ACP reductase transit peptides would also operate

20 effectively in plants such as rape embryo. The advantage of using the same plastid transit peptide for all three polyhydroxyalkanoate genes is to ensure that any variability in the uptake of the genes is not due to the transit peptide which is

25 used.

Although some proteins appear to be efficiently targeted to the plastid stroma by the transit peptide alone, other proteins also require the presence of up to twenty amino acids of the

30 amino terminus of the mature protein. The requirement for the presence of mature sequences appears to depend on the size and charge of the protein to be transported.

To obtain synthesis of polyhydroxyalkanoate

- 9 -

polymer in plant tissues it is necessary to obtain plants expressing all three genes for the enzymes β -ketothiolase, acetoacetyl-CoA reductase and polyhydroxybutyrate synthase. This may be achieved by using one of the following strategies:

- i) Plants may be individually transformed with the three polyhydroxyalkanoate pathway genes. Plants containing individual genes are grown up in the glass-house and cross-pollinated to obtain hybrid plants containing two pathway genes. This procedure is then repeated to produce hybrid plants containing all three genes.
- ii) Plants may be sequentially transformed with plasmids containing the individual pathway genes.
- iii) Two or three pathway genes may be cotransformed into the same plant by simultaneous infection with Agrobacteria containing the individual genes.
- iv) Plants may be transformed with plasmids containing two or three pathway genes.

A combination of these techniques may be used to obtain expression of all three genes in a single plant. Successive rounds of cross-pollination are carried out until the progeny are homozygous for all three genes. For methods (ii) and (iii) above, it is advantageous to insert each gene into vectors containing different selectable marker genes to facilitate selection of plants containing two or three polyhydroxyalkanoate pathway genes. Examples of selectable markers are genes conferring resistances to kanamycin, hygromycin, sulphonamides and bialaphos or phosphinothricin.

The invention will now be described by way of example only with reference to the accompanying drawings, in which:

5 Figure 1 shows the pathway for polyhydroxybutyrate production in Alcaligenes eutrophus;

Figure 2 is a physical map of the 5.2kb SmaI-EcoRI fragment of Alcaligenes eutrophus DNA;

10 Figure 3 is a map of the plant expression vector pJR11;

Figure 4 shows the Southern blot analysis of the three PCR products produced during the making of the ssu transit peptide-ketothiolase construct;

15 Figure 5 is a graph of β -ketothiolase enzyme activities in tobacco leaves;

Figure 6 is a graph of NADP acetoacetyl CoA reductase enzyme activities in tobacco leaves.

EXAMPLE

20 A 5.2kb SmaI-EcoRI fragment which codes for all three polyhydroxyalkanoate (PHA) biosynthetic genes had previously been isolated from Alcaligenes eutrophus (Schubert et al, 1988, J Bacteriol, 170). This fragment cloned into the vector pUC9 (New England Biolabs) together with a 2.3kb PstI sub
25 fragment cloned into Bluescript KS- (Stratagene) were provided by Dr Steinbuchel of the University of Gottingen, Germany. A restriction map of the fragment is shown in Figure 2. The positions of the restriction sites and the positions of the
30 genes for β -ketothiolase, acetoacetyl CoA reductase, and polyhydroxybutyrate (PHB) synthase are shown.

The expression vector chosen to gain constitutive expression of PHA biosynthetic genes

- 11 -

in tobacco and other rape plants was pJRli. This vector contains the cauliflower mosaic virus CamV35S promoter and the nos terminator, separated by a multiple cloning site to allow the insertion of the PHA genes. The vector also contains the kanamycin resistance nptII gene as a selectable marker. Figure 3 is a map of the plant expression vector pJRli. Vector pJRlRi was also utilised; this vector contains the expression cassette in the opposite orientation.

All routine molecular biological techniques were those of Sambrook et al (1989, A laboratory manual, Second edition). Oligonucleotides were all synthesised on an Applied Biosystems 380B DNA Synthesiser. PCR machines used were Techne PHC-1 Programmable Dri-Blocks. Taq polymerase was obtained from Perkin-Elmer/Cetus. Restriction enzymes and other modifying enzymes were obtained from New England Biolabs, Gibco/BRL, Northumbria Biologicals Limited and Pharmacia. Sequencing kits were obtained from Cambridge Biosciences (Sequenase) and Promega (Taqtrack). All radio-isotopes were supplied by Amersham International.

1. Construction of vectors to gain constitutive cytosolic expression of PHA pathway genes.

1.1. β -ketothiolase

The β -ketothiolase gene was isolated as a 1.3kb PstI-PleI fragment from the 2.3kb PstI fragment of pKS-::2.3P7. This fragment was blunt-ended with Klenow and was inserted into the dephosphorylated SmaI site of pJRli. The resulting plasmid was denoted pJRliT. Recombinant plasmids were identified by colony hybridisation using the

1.3kb ins rt fragment as a pr b . R striction
mapping of r combinants revealed those containing a
single β -ketothiolase insert in the sense
orientation. The orientation of the insert was
5 confirmed by sequencing using a primer that
hybridised to the 3' end of the CaMV35S promoter.

1.2. Acetyoacetyl-CoA reductase

The acetoacetyl-CoA reductase gene was
isolated as a 0.9kb *Ava*II-*Xmn*II fragment from pKS
10 ::2.3P7. This fragment was inserted into pJRIi as
described for pJRIiT. However, the orientation of
the insert fragment in recombinant plasmids could
not be confirmed by restriction mapping due to the
unavailability of suitable restriction enzyme
15 sites. Therefore four recombinants were sequenced
using the CaMV35S 3' primer and, of these, one was
found to contain a sense insert. This plasmid was
denoted pJRIiR.

1.3. PHB synthase

20 The PHB synthase gene was isolated from pKS-
::2.3P7 as a *Bst*BI-*Stu*I fragment. This fragment
was blunt-ended and inserted into pJRIi as
described for pJRIiT and pJRIiR. The identity of
recombinant (pJRIiS) plasmids containing a single
25 insert in the sense orientation was confirmed by
restriction mapping and by sequencing with the
CaMV35S 3' primer.

2. Construction of vectors for constitutive plastid targeted expression of PHA pathway enzymes.

30 Transport into plastids of the component
polypeptides for each of the PHB pathway enzymes
can be achieved by addition of a transit peptide
sequence to the 5' end of the gene sequence.

The first gene to be tailored was

ketothiolase. A technique involving polymerase chain reaction (PCR) was employed in order to join the p a RUBISCO small subunit transit p ptide sequence in frame with the ketothiolase gene.

5 Linking the transit peptide to the ketothiolase gene involved three experiments. The first experiment added a small portion of the 5' end of the ketothiolase gene onto the 3' end of the transit peptide sequence. The second experiment
10 added a small portion of the 3' end of the transit peptide onto the 5' end of ketothiolase gene. The third experiment utilised the overhangs produced in the preceding experiments to extend across the junction and produce full length transit peptide
15 linked in frame with the ketothiolase gene.

Four PCR primers were designed:-

1. 5' end of the transit peptide allowing extension toward its 3' end:

AAA TGG CTT CTA TGA TAT CCT CTT CAG CT

20

TP1

2. 3' end of transit peptide linked to 5' end of ketothiolase gene allowing extension toward 5' end of transit peptide:

ACG ATG ACA ACG TCA GTC ATG CAC TTT ACT CTT

25

CCA CCA TTG CTT GT

TPKB

3. 3' end of transit peptide linked to 5' end of ketothiolase gene allowing extension toward the 3' end of the ketothiolase gene:

30

ATT ACA AGC AAT GGT GGA AGA GTA AAG TGC

ATG ACT GAC GTT GTC ATC GT

TPKT

4. 3' end of ketothiolase gene:

ACC CCT TCC TTA TTT GCG CTC GAC T

K1

- 14 -

For the first experiment template DNA was pSM64 (transit peptide sequence) and the primers were TP1 and TPKB with an annealing temperature of 65°C. The derived PCR products were run out on an agarose gel and the band corresponding to 199bp cut out and electroeluted from the gel.

In the second experiment template DNA was pKS::2.3P7, the primers involved were TPKT and K1 and the annealing temperature 68°C. The products of the PCR reaction were again run out on a gel and the required 1.207kb band isolated and electroeluted from the gel slice.

The third experiment utilised the DNA isolated from the previous experiment as template and the primers TP1 and K1. The annealing temperature was 65°C and although this PCR experiment was very inefficient some full length product (1.352kb) was formed.

A small portion of each of the three PCR products was run out on an agarose gel. Southern blot analysis using three of the oligos as probes (TP1, K1 and TPKT) was carried out. Results are given in Figure 4 and show that the product of the third reaction contained the 5' end of the transit peptide, the overlap of 3' transit peptide and 5' ketothiolase gene, and the 3' end of the ketothiolase gene.

It was necessary to check the sequence of this product as it is known that PCR can incorporate base mismatches. The PCR product was blunt-ended and cloned into SmaI cut and phosphatased pUC18. Six clones were identified which contained the PCR product. The clones were sequenced using the universal and reverse primers (Sequenase kit and

Taqtrack kit). Clones with completely correct sequence through the transit peptide and the 5' end of the ketothiolase gene up to a TthIII1 restriction site within the gene were identified.

5 From one of these clones a TthIII1-Kpn1 fragment was excised. The Kpn1 site was cut back to give a blunt end, and a TthIII1-Sma1 fragment of Alcaligenes eutrophus DNA from pKS-::2.3P7 corresponding to the major portion of the
10 ketothiolase gene was inserted. Positive clones were sequenced across the joins. The transit peptide-ketothiolase fragment was excised and inserted into pJR1Ri.

For the transit peptide-reductase construct
15 PCR was also utilised. This required only one PCR experiment as a Dde I site (unique in the transit peptide and reductase sequences) was present close to the 5' end of the gene. The PCR experiment required two primers:-

20 1. Sequence homologous to the 5' end of the transit peptide which would allow extension toward the 3' end. A Cla I site was incorporated into the sequence 5' to the transit peptide sequence.

25 ACC ATC GAT GGA TGG CTT CTA TGA TAT CCT CTT
CAG CT

CLATP

2. Sequence homologous to just past the Dde I site in the reductase gene, linked in frame with 3' transit peptide sequence to allow
30 extension toward the 5' transit peptide.

ATG CGC TGA GTC ATG CAC TTT ACT CTT CCA CCA
TTG CTT GTA AT

TPDDER

After PCR with these two primers and transit peptid DNA as template the 195 bp product was identified on agarose gels and isolated by electroelution. DdeI XmnI reductase gene was isolated and ligated to DdeI cut PCR product. After agarose gel electrophoresis the 1.063 kb band was isolated, cut with ClaI and ligated into ClaI EcoRV Bluescript SK(-). Positives are being characterised.

3. Transformation of plants with the PHB genes

3.1. Agrobacterium transformations

Cesium-pure pJRIiT, pJRIiR, pJRIiS and pJRIi were individually transformed into Agrobacterium tumefaciens strain LBA4404 by direct uptake as follows. LB (10mls) was inoculated with A tumefaciens strain LBA4404. The culture was shake-incubated at 28°C for approximately 16 hours until the optical density (OD) at 660 nm was 0.5. The cells were recovered by centrifugation (3000 rpm Sorvall RT6000B, 6 mins, 4°C). They were resuspended in 250µl of ice-cold 20mM CaCl₂. The cell suspension was then dispensed into pre-chilled Eppendorf tubes in 0.1ml aliquots. Approximately 1µg of caesium-pure plasmid DNA was added to each tube. The cells were then heat-shocked by freezing in liquid nitrogen followed by incubation at 37°C for 5 minutes. LB medium (1ml) was added and the cells were allowed to recover by incubation (shaken) at 28°C for 3-4 hours. The cell pellets were obtained by centrifugation (11,500g, 30 seconds, 20°C) and resuspended in 0.1ml LB. Recombinant cells were selected on LB (agar-solidified) containing kanamycin (50µg/ml), streptomycin (500µg/ml) and rifampicin (100µg/ml)

following incubation at 28°C. Mini-prep DNA of the resultant Agrobacterium strains was then isolated and analysed by restriction enzyme digestion to ensure that no re-arrangements had occurred.

5 **3.2. Plant Transformations**

Tobacco leaf pieces and oilseed rape petioles were inoculated individually with strains LBA4404/JRIi, LBA4404/pJRIiT, LBA4404/pJRIiR and LBA4404/pJRIiS. Plants were cultured in a growth
10 room with a temperature of 25°C and a photoperiod of 16 hours.

Brassica napus cv. Westar seedlings were sterilised in 10% sodium hypochlorite and washed in sterile water before germination on MS medium
15 (Imperial)(containing 3% sucrose and 0.7% phytagar (Gibco). The cotyledons were excised from 5 day old seedlings and the petioles of which were placed in MS medium as above but supplemented with 4.5µg/ml benzylaminopurine (BAP). The cotyledons
20 were cultured in this medium for 24 hours after which their petioles were dipped in an Agrobacterium solution. The Agrobacterium culture had been grown overnight in LB medium containing kanamycin (50µg/ml) following which the
25 Agrobacterium cells had been pelleted and washed in liquid MS medium and diluted to OD₆₆₀ 0.1. The inoculated petioles were returned to the MS medium containing 4.5µg/ml BAP and incubated in the culture room for 2 days. The cotyledons were then
30 transferred to MS medium supplemented with BAP (4.5µg/ml), carbenicillin (Duchefa) (500µg/ml) and kanamycin (15µg/ml). The cotyledons were subcultured on this medium every 2 weeks until the production of green callus and eventually shoots.

Shoots were excised and cultured on MS containing carbamycin (500 µg/ml) and kanamycin (15 µg/ml) until they were transferred to the glasshouse.

5 Nicotiana tabacum cv SRI seeds were sterilised as described above and germinated on MS medium (containing 3% sucrose and 0.8% bactoagar). The shoot tips from these seedlings were then micropropagated on this media to provide plants for transformation studies. Leaf pieces from these
10 plants were dipped in an Agrobacterium solution (prepared as described above) and were then cultured on MS medium containing 3% sucrose, 0.8% bactoagar, 1 µg/ml BAP and 0.1 µg/ml NAA, for 2 days. The leaf pieces were then cultured on the same
15 media supplemented with carbenicillin (500 µg/ml) and kanamycin (100 µg/ml) for 5 weeks. Regenerated shoots were excised and cultured on MS containing 3% sucrose, 0.8% bactoagar, 200 µg/ml carbenicillin and 100 µg/ml kanamycin for 2 passages of 5 weeks
20 before transfer to the glasshouse.

Kanamycin-resistant tobacco and rape plants were obtained for those transformed individually with JRIi, JRIiT, JRIiR and JRIiS.

3.3. Cotransformations

25 Rape cotyledons and tobacco leaf pieces were also inoculated with mixtures of Agrobacterium strains. These inoculations were performed as described previously except that 1:1 mixtures of diluted Agrobacterium cultures, of the same optical
30 density, were prepared immediately prior to inoculation.

4. Biochemical assessment of plants

Expression of Alcaligenes eutrophus PHA pathway enzymes in plant tissues was detected by

- 19 -

nzyme activity assays. The presence of the enzyme polypeptides was also detected by Western blot analysis.

For the latter analyses rabbit polyclonal antibodies were raised to the purified β -ketothiolase and NADP acetoacetyl CoA reductase enzymes from Alcaligenes eutrophus. Bacteria were pelleted, washed, and crude extracts prepared as described by Haywood and Large (1981, Biochem J, 199, 187-201). β -ketothiolase A was purified by chromatography on hydroxylapatite, followed by anion exchange chromatography on FPLC mono Q, followed by gel filtration on Superdex S-200 (Pharmacia), using modifications of methods described by Haywood et al (1988, FEMS Microbiology Letters, 52, 91-96). NADP acetoacetyl-CoA reductase was purified using the same techniques, with an additional affinity chromatography step on 2',5' ADP sepharose (Pharmacia). Purified proteins were subjected to sodium dodecyl sulphate polyacrylamide gel electrophoresis (SDS PAGE) according to the method of Laemmli (1970, Nature, 222, 680-685). The final β -ketothiolase preparation showed a single coomassie blue stained band at 41kd. The final reductase preparation showed a major band at 26kd. 3mg of purified ketothiolase and 2mg of purified reductase were subjected to preparative SDS PAGE. The bands corresponding to the two enzymes were electroeluted from the gels and injected into rabbits to raise polyclonal antibodies. Sera from primary and secondary bleeds following injection were shown to contain antibodies specific for their target enzymes via Western blot analyses of crude

Alcaligenes extracts.

Crude extracts of tobacco leaves were prepared by grinding leaf tissue in 50mM potassium phosphate buffer pH7.0 containing 1mM dithiothreitol. After centrifugation at 30,000g, enzyme assays for ketothiolase and acetoacetyl CoA reductase were conducted on aliquots of the supernatants by the methods described by Haywood et al (1988, FEMS Microbiology Letters, 52, 91-96; 52, 259-264). PHB synthase assays were conducted on aliquots of the 30,000g supernatants and aliquots of the pellets, resuspended in extraction buffer, by the method of Haywood et al (1989, FEMS Microbiology Letters, 57, 1-6).

For Western blot analysis, aliquots of the 30,000g supernatants were subjected to SDS PAGE and electrophoretically transferred to nitrocellulose filters. Filters were then rinsed in TBS (50mM Tris-HCl pH7.9, 150mM NaCl) and incubated in TBS plus 5% bovine serum albumin. Proteins reacting with anti-ketothiolase or anti-reductase serum were detected by incubating the filters in 100ml TBS containing 2ml of the relevant serum for 1-2h. Bound first antibody was subsequently detected using goat anti-rabbit IgG alkaline phosphatase conjugate and nitroblue tetrazolium alkaline phosphatase colour development reagent (BioRad Laboratories).

Initial biochemical analyses were carried out on subcultured tobacco plants growing in tissue culture. Eighteen kanamycin resistant plants transformed with JR11 ketothiolase were subjected to enzyme analysis and results compared with untransformed control plants. Leaves of the same

- 21 -

size were extracted.

Figure 5 shows the β -k tothiolase nzym activities in the tobacco l aves. The identification numbers of individual plants are shown on the x axis. Plants to the left of the dotted line are untransformed control plants. Plants to the right of the line are transformed with JR1i ketothiolase.

A low level of ketothiolase activity was detected in untransformed control plants. Nearly all of the JR1i ketothiolase transformed plants had ketothiolase activity higher than control. The highest activity was 34 nmol/min/mg protein, 2.8 times higher than the highest control plant. In Western blots the anti-ketothiolase antibody detected a polypeptide at 41kd in untransformed control tobacco plants - possibly corresponding to the endogenous ketothiolase enzyme activity. While a 41kd polypeptide was also detected in extracts of JR1i ketothiolase transformed plants, the Western blots could not quantitatively distinguish transformed from untransformed plants.

Figure 6 shows the NADP acetoacetyl CoA reductase enzyme activities in leaves of the tissue culture grown tobacco plants. The identification numbers of individual plants are shown on the x axis. Plants to the left of the dotted line are untransformed control plants. Plants to the right of the line are transformed with pJR1i reductase.

A low level of acetoacetyl CoA reductase activity was detected in untransformed control plants. Nearly all the 21 JR1i reductase transformed plants had reductase activity higher than control. The highest activity was 30

- 22 -

nmol/min/mg protein, 4 fold higher than the high st control plant. In Western blots the anti-reductase antibody did not detect any polypeptide with a m.w. of 26kd in extracts of untransformed control plants. A 26kd polypeptide was however detected in extracts of the JR1i reductase transformed plants. Expression of the bacterial reductase gene in tobacco leaves was therefore demonstrated.


CLAIMS

1. A plant adapted for the production of polyhydroxyalkanoate comprising a recombinant genome of an oil-producing plant, which genome contains genes encoding enzymes necessary for catalysing the production of polyhydroxy-alkanoate together with gene regulatory sequences directing expression of the said genes to target plant cell components.
2. A plant according to claim 1 wherein the genes encoding the enzyme or enzymes necessary for the catalysis of polyhydroxyalkanoate production are isolated from a micro-organism.
3. A plant according to claim 2 wherein the micro-organism is Alcaligenes eutrophus.
4. A plant according to claim 1 in which then said oil-producing plant is selected from the group consisting of oilseed rape, canola, soya and sunflower.
5. A plant as claimed in claim 1, in which the said enzyme is selected from the group consisting of β -ketothiolase, acetoacetyl-CoA reductase and polyhydroxybutyrate synthase.

6. A plant as claimed in claim 1, in which the said gene regulatory sequences direct expression of the polyhydroxyalkanoate genes to the oil storage organ.
7. A plant as claimed in claim 6, in which expression is directed to the developing oil storage organ.
8. A plant as claimed in claim 6, in which expression is directed to the embryo.
9. A plant as claimed in claim 1, in which the said gene regulatory sequences direct expression of the polyhydroxyalkanoate genes to the cytosol or to the mitochondrion or to the plastid.


1 / 7

FIG. 1



2 / 7

FIG. 2



3/7

Restriction enzyme sites.

AccI	73, 1928, 2066, 3116, 3131
AvaI	2, 1336, 2964, 3438
AvaII	751, 944, 1126, 1602, 2232, 2297, 2495, 3614, 3630, 3915
BamHI	-
BanII	-
BglII	1424
BspMI	661, 1211, 1613, 2114, 2144, 2171, 2186, 2652, 2826, 4307, 4703
DdeI	176, 1080, 3958, 4898
EcoRI	5196
HincII	-
HindIII	-
KpnI	2163, 2200
NcoI	1097, 4444
NotI	983
PleI	527, 1368, 2442, 3950, 4143, 4643
PstI	1610, 2183, 2663, 4984
Sau3AI	-
SacI(SstI)	1621, 2192
Sall	1927, 2065, 3115, 3130
SmaI	4
SphI	-
StuI	2657, 3652
XbaI	-
XmaI	-
XmnI	1075, 4826

FIG. 2 (2/2)

4 / 7





FIG. 3 The plant expression vector pJR1i

5/7


FIG. 4



- M markers
1 PCR product from experiment 1
2 PCR product from experiment 2
3 PCR product from experiment 3


6/7

FIG. 5



7/7


FIG.6



INTERNATIONAL SEARCH REPORT

International Application

PCT/GB 92/00742

I. CLASSIFICATION OF SUBJECT MATTER (If several classification symbols apply, indicate all) ⁶		
According to International Patent Classification (IPC) or to both National Classification and IPC		
Int.Cl. 5 C12N15/82; C12N15/52; A01H5/00		
II. FIELDS SEARCHED		
Minimum Documentation Searched ⁷		
Classification System	Classification Symbols	
Int.Cl. 5	C12N ; C12P ; A01H	
Documentation Searched other than Minimum Documentation to the Extent that such Documents are Included in the Fields Searched ⁸		
III. DOCUMENTS CONSIDERED TO BE RELEVANT⁹		
Category ¹⁰	Citation of Document, ¹¹ with indication, where appropriate, of the relevant passages ¹²	Relevant to Claim No. ¹³
Y	WO,A,9 100 917 (MIT) 24 January 1991 see page 45, line 24 - page 47, line 30 ---	1-9
Y	EP,A,0 189 707 (PLANT GENETIC SYSTEMS) 6 August 1986 see the whole document ---	1-9
A	JOURNAL OF BIOLOGICAL CHEMISTRY. vol. 264, no. 26, 15 September 1989, BALTIMORE US pages 15298 - 15303; PEOPLES, O.P., ET AL.: 'Poly-beta-hydroxybutyrate (PHB) biosynthesis in Alcaligenes eutrophus H16' see figure 4 --- -/-	1-9
<p>¹⁰ Special categories of cited documents:</p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>"&" document member of the same patent family</p>		
IV. CERTIFICATION		
Date of the Actual Completion of the International Search	Date of Mailing of this International Search Report	
2 18 AUGUST 1992	01. 09. 92	
International Searching Authority	Signature of Authorized Officer	
EUROPEAN PATENT OFFICE	MADDOX A.D. 	

III. DOCUMENTS CONSIDERED TO BE RELEVANT (CONTINUED FROM THE SECOND SHEET)		Relevant to Claim No.
Category *	Citation of Document, with indication, where appropriate, of the relevant passages	
A	<p>SCIENCE. vol. 245, 15 September 1989, LANCASTER, PA US pages 1187 - 1189; POOL, R., ET AL.: 'In search of the plastic potato' see page 1189, column 2 - column 3 ---</p>	1-9
A	<p>J. BACTERIOLOGY vol. 170, no. 12, December 1988, pages 5837 - 5847; SCHUBERT, P., ET AL.: 'Cloning of the Alcaligenes eutrophus genes for the synthesis of poly-beta-hydroxybutyric acid (PHB) and synthesis of PHB in Escherichia coli' cited in the application see the whole document ---</p>	1-9
A	<p>BIOLOGICAL ABSTRACTS vol. 82, (1986) ref. no. 86881 & PLANT PHYSIOLOGY. vol. 81, no. 3, 1986, ROCKVILLE, MD, USA. pages 817 - 822; BOYLE, S.A., ET AL.: 'Uptake and processing of the precursor to the small subunit of ribulose-1,5-bisphosphate carboxylase by leucoplasts from the endosperm of developing castor oil (Ricinus communis cultivar Baker 296) seeds' see the abstract ---</p>	1-9

**ANNEX TO THE INTERNATIONAL SEARCH REPORT
ON INTERNATIONAL PATENT APPLICATION NO.**

GB 9200742
SA 58672

This annex lists the patent family members relating to the patent documents cited in the above-mentioned international search report.
The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information. 18/08/92

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO-A-9100917	24-01-91	EP-A- 0482077	29-04-92
EP-A-0189707	06-08-86	AU-B- 591087	30-11-89
		AU-A- 5166485	17-07-86
		JP-A- 61224990	06-10-86